

Variabilitätsfaktoren für die Abschätzung der Kurzzeitexposition von Verbrauchern gegenüber Pflanzenschutzmittelrückständen

Information Nr. 014/2010 des BfR vom 5. März 2010

In Studien, bei denen geprüft wird, wie hoch der Rückstand eines Pflanzenschutzmittel-Wirkstoffs nach ordnungsgemäßer Anwendung im Ernteprodukt ist, werden normalerweise nicht Rückstände in einzelnen Einheiten, sondern in Mischproben aus mehreren Einheiten bestimmt. So erhält man ein insgesamt repräsentativeres Bild der Rückstandssituation. Sofern nicht von einer gleichmäßigen Vermischung einzelner Einheiten des Lebensmittels vor dem Verzehr auszugehen ist (z.B. Getreide), muss immer der Fall in der Bewertung berücksichtigt werden, dass Verbraucher eine einzelne, hoch belastete Einheit eines Lebensmittels verzehren. Der kritischste, aber nicht auszuschließende Fall ist demnach, dass 1. der in einer Mischprobe (z.B. bestehend aus einem Kilo Äpfel) gemessene Rückstand ausschließlich auf hohe Rückstände einer einzigen Einheit (z.B. eines Apfels) zurückgeht, während die übrigen Einheiten alle rückstandsfrei waren, und dass 2. dieser eine Apfel vom Verbraucher verzehrt wird. Um sicherzustellen, dass Mischproben nicht zu einer Unterschätzung eines möglichen akuten Risikos führen, wird der Rückstand in der Mischprobe daher mit einem Variabilitätsfaktor multipliziert. Standardmäßig werden Variabilitätsfaktoren von 5 bzw. 7 (in Abhängigkeit vom Gewicht einer Einheit) verwendet.

Auf der Basis von empirischen Daten aus Rückstandsstudien mit Pflanzenschutzmitteln, bei denen einzelne Einheiten anstelle von Mischproben untersucht wurden („Variabilitätsstudien“), wurde jedoch eine Reihe spezifischer Variabilitätsfaktoren abgeleitet und im Pflanzenschutzmittel-Zulassungsverfahren oder bei der Festsetzung von Rückstandshöchstgehalten für Pflanzenschutzmittel-Wirkstoffe verwendet. Die spezifischen Faktoren liegen in den meisten Fällen zwischen 2 und 3 und ersetzen für die jeweilige Wirkstoff/Lebensmittel-Kombination den Standard-Variabilitätsfaktor.

Das Bundesinstitut für Risikobewertung (BfR) stellt Informationen zu diesen spezifischen Variabilitätsfaktoren zur Verfügung und diskutiert kritisch, unter welchen Bedingungen sie für die Risikobewertung verwendet werden können. Das Dokument wurde in englischer Sprache verfasst.

1 Summary

Based on empiric data from unit-to-unit variability studies a couple of variability factors, which deviate from the normally applied default values, were derived for the evaluation of supervised field trials data in the context of registration and MRL setting procedures for pesticides. In addition to these special unit-to-unit studies, market surveys were available trying to estimate the variability within lots which have probably not been subject to mixing or blending. Since the true nature of these lots is, however, unknown, they were not used to derive variability factors from.

The variability factors from supervised unit-to-unit variability studies were in most cases between 2 and 3 and were mostly derived for specific pesticide/commodity combinations. They do not necessarily apply for residues of the same pesticide in other commodities or for residues of similar pesticides in the same commodity.

It should be noted, that

- no guideline is available yet on how to conduct unit-to-unit variability studies;

- no consensus exists yet concerning the adequate accuracy of the variability factor (BfR prefers to round the values to integral numbers and use those in the assessment);
- not all of the factors mentioned herein have been confirmed on EU level and have been used to base authorization/MRL decisions on;
- the derived factors are only applicable to lots clearly identified as unblended or unmixed.

For the acute exposure assessment of market samples without clear evidence of dealing with unblended lots the default factors of 5 for units above 250 g and of 7 for units between 25 and 250 g should be applied.

2 Assessment

2.1 Introduction

Samples taken from supervised field trials are normally collected randomly and are then combined, homogenized and analyzed as a composite sample. Variability factors are used in the acute dietary intake assessment of pesticides to account for the unit-to-unit variability in composite samples, i.e. to account for the situation, in which the whole residue measured in the composite sample originated from one single unit only. Consumption of such a “high residue unit” may occur in real life and needs to be considered in risk assessment as a reasonable worst case.

2.2 Acute dietary intake assessment and the use of default variability factors

The acute dietary intake of pesticide residues is calculated using the so-called IESTI equation (IESTI = International Estimated Short-Term Intake). Four different versions of the equation are used for different types of commodities. Case 1 is the simple case where the residue in a composite sample reflects the residue level in a meal-sized portion of the commodity (unit weight < 25 g, i.e. cherries). Case 2 is the situation where the meal-sized portion as a single fruit or vegetable unit might have a higher residue than the composite sample. Case 2 is further divided into case 2a and case 2b where the unit size is less than or greater than the large portion size, respectively. Case 3 allows for the likely bulk-ing and blending of processed commodities such as flour, vegetable oils and fruit juices. For more details the reader is referred to the EFSA opinion on acute dietary intake assessment (EFSA, 2007) and to the FAO manual (FAO, 2009).

The two versions of the IESTI equation in which variability factors play an important role, namely case 2a and case 2b, are depicted below:

$$\text{Case 2a} \quad IESTI = \frac{U \times (HR \text{ or } HR - P) \times v + (LP - U) \times (HR \text{ or } HR - P)}{bw}$$

$$\text{Case 2b} \quad IESTI = \frac{LP \times (HR \text{ or } HR - P) \times v}{bw}$$

U	Unit weight of the commodity
HR or HR-P	Highest residue or Highest residue after processing
v	Variability factor
LP	Large portion
bw	Body weight

According to the International Conference on Pesticide Residues Variability and Acute Dietary Risk Assessment held in York, UK (Harris et al., 2000), the variability factor is defined as the ratio between the 97.5th percentile and the mean of residues in individual units:

$$v = \frac{97.5th\ Percentile}{Mean}$$

In the frame of authorisation of plant protection products, the HR is estimated on the basis of supervised field residue trials in which the plants received treatments according to the most critical intended conditions of use (called “critical GAP”). As described in the introduction, composite samples of randomly selected units are usually considered to be representative of the mean residue in commodities from one treated field. To account for the theoretical 97.5th percentile present in the treated field and to cover a single high residue present in a single unit, the residue in the composite sample has to be multiplied with the variability factor.

Standard variability factors as included in EFSA PRIMo¹ are 5 (unit weight > 250 g) and 7 (unit weight between 25 g and 250 g). These are normally used within the EU for risk assessment and MRL setting purposes. For head lettuce and head cabbage, the standard variability factor is ambiguous. Factors 3 and 5 are currently applied. In its position paper for the 35th session of the CCPR in 2003 (EU Commission, 2003) the EU recommended to use a default variability factor of 3 for head lettuce and head cabbage. Consequent to this decision a factor of 3 was also used by PPR Panel in its opinion concerning the impact of a possible change of variability factors on the overall level of protection (EFSA, 2007). In the current version of EFSA PRIMo a default factor of 5 is normally pre-set for head lettuce and head cabbage. The guidance given on notification criteria for pesticide residue findings in the RASFF system (EU Commission, 2004) is still utilizing the variability factors of 5 (unit weight > 250 g) and 7 (unit weight ≤ 250 g), irrespective of the commodity concerned.

It has to be noted, that the IESTI equation is currently under revision. Chapter 2.2 refers to the currently applied approach.

2.3 Acute dietary intake assessment and the use of specific variability factors

BfR has been provided with unit-to-unit variability studies for a couple of pesticide/commodity combinations. Some investigations were related to classes of active substances rather than to single pesticides. In addition, information is available from published literature. EFSA also reported a couple of additional data from both supervised trials and market surveys (EFSA, 2005), which have not been available to BfR and are therefore not reported herein.

The available study results are summarized in detail in the appendix to this document. All studies reported (apart from chlorpropham, which was applied post-harvest) comprise at least analyses of residues in 100 single units. Hamilton et al. (Hamilton et al., 2004) showed that a sample size of 119 or more commodity units is required to achieve a 95 % certainty that at least one unit exceeds the 97.5th percentile of the sampled population. In its opinion on appropriate variability factors (EFSA, 2005), the PPR Panel considered to take also data-sets with at least 50 individual residue values into account for the estimation of the variability.

In the following table, variability factors derived from supervised unit-to-unit variability studies are summarized. If more than one study was available for one pesticide/commodity combination, the mean variability factor is given. Results from market surveys are presented in the

¹ http://www.efsa.europa.eu/EFSA/General/calculation_acutechronic_rev2.xls?ssbinary=true

overview table at the end of this document (appendix), but were not used to derive variability factors from.

Commodity	Pesticide	Variability factor
Apples	Azinphos-methyl	2.4
	Captan	2.7
	Dithianon	2.9
	Fenpyroxymate	2.2
	Pyraclostrobin	2.6
	Imazalil	1.5 (post-harvest treatment only)*
Grapes	<i>Anilinopyrimidine</i> **	2.6**
	<i>Dicarboximide</i> **	3.4**
	<i>Organophosphate</i> **	2.6**
	Pyraclostrobin	2.3
	<i>Pyrethroid</i> **	2.5**
	<i>Triazole (group)</i> **	2.5**
Potatoes (stored)	Chlorpropham	3.1
Head cabbage	all pesticides	3***
Head lettuce	all pesticides	3***

* This value is only applicable, if the treatment is known to have occurred post harvest

** One active substance from the whole group has been tested, but its identity has not been reported. It is therefore questionable, if the derived variability factor can be used for all compounds belonging to the respective group. No international agreement has been reached concerning the use of the mentioned factors. While for head cabbage and lettuce the use of an overall variability factor of 3 has been supported by JMPR and the EU Commission, this is not the case for grapes. BfR therefore does not recommend to use the mentioned factors for the time being.

*** Since only pesticide groups instead of specific pesticidal substances have been reported (see also the remark above), not the individually derived factors (< 3 each) but a slightly higher overall factor of 3 was recommended by the JMPR for head cabbage and lettuce. This accounts for uncertainties when extrapolating to all compounds belonging to the respective group.

2.4 Applicability of specific variability factors

The specific variability factors mentioned above have been derived for use together with supervised field trials data in the context of registration and MRL setting procedures.

Acute consumer risk assessment in the context of evaluating monitoring/surveillance samples is principally conducted as described in chapter 2.2, i.e. using the IESTI equation. The HR is, however, replaced by the OR (observed residue). OR is the residue concentration which has been measured in the respective market sample. In contrast to samples from supervised field trials, market samples often consist of different lots which have been mixed prior to marketing. This mixing may result in an increase of the variability, since treated and untreated lots may have been blended thus resulting in a lower overall mean residue while high residues on individual units remain unchanged.

In its opinion related to the appropriate variability factors to be used for acute dietary exposure assessment (EFSA, 2005), the PPR Panel confirmed that “variability factors estimated from samples collected in the marketplace were higher than those from samples obtained in experimental studies (supervised trials).” The PPR Panel therefore recommended that consideration should be given to using different variability factors when doing exposure assessments with data from supervised trials and from monitoring/surveillance. BfR and BVL have published an opinion on this issue in 2007 (Banasiak et al., 2007) and came to the conclusion that as long as standard variability factors of 5 and 7 are used in exposure assessments with data from supervised trials, the same factors may be used and are still considered protective for the assessment of monitoring/surveillance samples. As mentioned in chapter 2.2,

the IESTI equation together with the variability factors in place is currently under revision. When the revision has been finalized, the recommendation as how to use the same standard variability factors for both assessment purposes will need to be revisited.

The situation is somewhat different concerning the use of specific variability factors. As set out in chapter 2.3, these factors are usually between 2 and 3. These factors might be applied in exposure assessments with data from supervised trials. For market surveys, however, distribution ranges of variability factors were found to be much broader (EFSA, 2005). In general the BfR therefore recommends applying lowered variability factors for pesticide/commodity combinations only in those situations, where the specific lot can be clearly identified as being unblended and evenly treated over all single units. A general use of lowered variability factors might result in an underestimation of the dietary intake and is not acceptable in terms of consumer health protection.

3 References

Banasiak et al., 2007: Banasiak, U., Herrmann, M., Hohgardt, K., Michalski, B. and Sieke, C., Abschätzung des akuten Risikos durch Pflanzenschutzmittel-Rückstände in Lebensmitteln auf der Basis von Daten aus amtlicher Überwachung und Eigenkontrollen (in German language with English summary), J. Verbr. Lebensm. 2 (2007), 54-60

Blaschke, U., 2007: Residue Study (Unit to Unit variation) with BAS 584 00 F applied to apples in Germany and Spain in 2006, 14.03.2007, BASF DocID 2007/1018308

Bross, M., 2006: Dithianon (BAS 216 F) - Supplemental information on supervised field trials – Study design and preliminary results of the unit-to-unit variability study in apples, 13.10.2006, BASF Doc ID 2006/1034351

EFSA, 2005: Opinion of the Scientific Panel on Plant health, Plant protection products and their Residues on a request from Commission related to the appropriate variability factor(s) to be used for acute dietary exposure assessment of pesticide residues in fruit and vegetables, The EFSA Journal (2005), 177, 1-61

EFSA, 2007: Opinion of the Scientific Panel on Plant protection products and their Residues on a request from the Commission on acute dietary intake assessment of pesticide residues in fruit and vegetables (Question N° EFSA-Q-2006-114), adopted on 19 April 2007, The EFSA Journal (2007) 538, 1-88, <http://www.efsa.europa.eu/en/scdocs/doc/538,0.pdf>

European Commission, 2003: European Community Position for the 35th Session of the Codex Committee on Pesticide Residues, Rotterdam, 31 March-5 April 2003, Point 2.9, http://ec.europa.eu/food/fs/ifsi/eupositions/ccpr/ccpr_ec-comments_35th_en.pdf

European Commission, 2004: Draft Guidance Document on Notification criteria for pesticide residue findings to the Rapid Alert System for Food and Feed (RASFF), SANCO/3346/2001 rev 7, Brussels, 14.07.2004
http://ec.europa.eu/food/plant/protection/resources/rasff_pest_res_en.pdf

FAO, 2009: FAO manual on the submission and evaluation of pesticide residues data for the estimation of maximum residue levels in food and feed, Second Edition, Rome, 2009
http://www.fao.org/fileadmin/templates/agphome/documents/Pests_Pesticides/JMPR/FAO_manual2nded_Oct07.pdf

Hamilton et al., 2004: Hamilton, D., Ambrus, A., Dieterle, R., Felsot, A., Harris, C., Petersen, B., Racke, K., Wong, S., Gonzalez, R., Tanaka, K., Earl, M., Roberts, G. and Bhula, R., Pesticide residues in food – acute dietary exposure, *Pest Manag Sci* 60 (2004), 311-339

Harris, C., 1998: Unit to Unit Variability of Pesticide Residues in Celery, Plums and kiwi fruit, Report by the Pesticides Safety Directorate, York, 1998

Harris et al, 2000: Harris, C.A., Mascall, J.M., Warren, S.F.P., Crossley, S.J., Summary report of the International Conference on pesticide residues variability and acute dietary risk assessment, *Food Additives and Contaminants* 17(7) (2000), 481-485

Harrison, C., 2007: Final Report on Project AF/10969/MK to determine the magnitude and variability of azinphos-methyl residues in individual fruit and the raw agricultural commodity apple resulting from three application of GUSATHION XL (Southern Europe, 2006), 26.02.2007; AF/10969/MK

Harrison, C., 2007a: Final Report on Project AF/10970/MK to determine the magnitude and variability of azinphos-methyl residues in individual fruit and the raw agricultural commodity apple resulting from three applications of GUSATHION XL (Southern Europe, 2006), 26.02.2007; AF/10970/MK

Heck, W., 2002: Determination of the Residue Variability of BAS 500 F and KIF-230 in grapes after treatment with BAS 525 00 F under field conditions in Germany and Spain, 2000, 30.10.2002, BASF DocID 2002/1008790

Kaethner, M., 2002: Determination of Residue Variability in Table and Wine grapes after a Tank-Mix Application of Anilinopyrimidine, Triazole, Pyrethroid, Organophosphate, and Dicarboximide Crop Protection Products – France/Germany 2000 to 2001, Summary Report by the ECPA Residues Expert Group, 2002, BASF DocID 2002/1007077

Kaethner, M., 2002a: Determination of Residue Variability in Head Lettuce following a Tank-Mix Application of Anilinopyrimidine, Triazole, Pyrethroid, Organophosphate, Carbamate, and Dicarboximide Crop Protection Products – France/Germany 2000 to 2001, Summary Report by the ECPA Residues Expert Group, 2002, BASF DocID 2002/1007078

Klimmek, S., 2006: Residue Variability of Captan and its Metabolite Tetrahydrophthalimide (THPI) in Individual Apple Fruit after Application of Merpan 80 WGD to a Commercial Apple Orchard with a Common Spraying Technique – Study Representative for the Fruit Group – Germany, Season 2005, 23.01.2006, Report No. MAC0501

Klimmek, S., Klimmek, A., 2007: Residue Variability of Fenpyroximate in Individual Apple Fruit after Application of Kiron 5% SC to a Commercial Apple Orchard with Common Spraying Technique – Study Representative for the Fruit Group – Germany, Season 2006, 05.03.2007; Report No. NNC-0601

Quirijns, J.K., 2003: Distribution of chlorpropham on individual potatoes in samples from bulk stores, 28.05.2003, TNO Report No. V5092

Tetuàn, B., 2007: Residue Behaviour and Variability of Residues of Imazalil after Post-Harvest Application of an Imazalil-based Formulation (UB027180) on Apple Fruits, Spain 2007, 29.06.2007, Report No. 20077006/S1-FPH

Appendix

Overview of unit-to-unit variability studies available to the BfR

Commodity	Pesticide	Number of units	Estimated variability factor	Type of survey	Remarks	Reference	
Kiwi fruit	Phosmet	100	6 ^a	Market sample ^c		Harris, C., 1998	
	Methidathion Parathion-methyl	100	5 ^a 2 ^a	Market sample ^c		Harris, C., 1998	
	Methidathion Parathion-methyl	100	4 ^a 2 ^a	Market sample ^c		Harris, C., 1998	
	Fenitrothion Quinalphos	100	Not possible ^b 5 ^a	Market sample ^c		Harris, C., 1998	
	Diazinon	100	3 ^a	Market sample ^c		Harris, C., 1998	
	Range (total mean)			2 – 6 (3.9)	Market sample^c		
Plums	Chlorpyrifos Pirimicarb	100	6 ^a Not possible ^b	Market sample ^c		Harris, C., 1998	
	Chlorpyrifos Phosalone	100	Not possible ^b Not possible ^b	Market sample ^c		Harris, C., 1998	
	Phosalone Pirimicarb	100	14 ^a 16 ^a	Market sample ^c		Harris, C. 1998	
	Pirimicarb	100	6 ^a	Market sample ^c		Harris, C., 1998	
	Phosalone	100	5 ^a	Market sample ^c		Harris, C., 1998	
	Acephate Methamidophos	100	4 ^a 4 ^a	Market sample ^c		Harris, C., 1998	
	Chlorpyrifos Dimethoate Omethoate	100	Not possible ^b 23 ^a Not possible ^b	Market sample ^c		Harris, C., 1998	
	Acephate Methamidophos Pirimiphos-methyl	100	3 ^a 4 ^a 8 ^a	Market sample ^c		Harris, C., 1998	
	Fenitrothion	100	5 ^a	Market sample ^c		Harris, C., 1998	
	Acephate Methamidophos	100	5 ^a 4 ^a	Market sample ^c		Harris, C., 1998	
	Range (total mean)			3 – 23 (7.6)	Market sample^c		

Commodity	Pesticide	Number of units	Estimated variability factor	Type of survey	Remarks	Reference
Apples	Dithianon	150	3.8	Supervised field trial		Bross, M., 2006
	Dithianon	150	2.6	Supervised field trial		Bross, M., 2006
	Pyraclostrobin Dithianon	150	2.7 2.4	Supervised field trial		Blaschke, U., 2007
	Pyraclostrobin Dithianon	150	2.4 1.9	Supervised field trial		Blaschke, U., 2007
	Pyraclostrobin Dithianon	150	2.7 3.8	Supervised field trial		Blaschke, U., 2007
	Pyraclostrobin Dithianon	150	2.7 2.6	Supervised field trial		Blaschke, U., 2007
	Azinphos-methyl	100	2.4	Supervised field trial		Harrison, C., 2007
	Azinphos-methyl	100	2.3	Supervised field trial		Harrison, C., 2007a
	Captan	100	2.7	Supervised field trial		Klimmek, S., 2006
	Fenpyroximate	130	2.2	Supervised field trial		Klimmek, S., 2007
	Range (total mean)		1.9-3.8 (2.7)	Supervised field trial		
Apples (post harvest)	Imazalil	119	1.5	Supervised post-harvest treatment		Tetuàn, B., 2007
		Range (total mean)	1.5	Supervised post-harvest treatment		
Grapes	Anilinopyrimidine Dicarboximide Organophosphate Pyrethroid Triazole	120	2.6 2.9 2.3 2.6 3.4	Supervised field trial	Individual compounds not specified	Kaethner, M., 2002
	Anilinopyrimidine Dicarboximide Organophosphate Pyrethroid Triazole	120	2.4 2.3 2.3 2.2 2.0	Supervised field trial	Individual compounds not specified	Kaethner, M., 2002
	Anilinopyrimidine Dicarboximide Organophosphate Pyrethroid Triazole	120	2.5 2.7 2.5 2.8 2.4	Supervised field trial	Individual compounds not specified	Kaethner, M., 2002

Commodity	Pesticide	Number of units	Estimated variability factor	Type of survey	Remarks	Reference
Grapes (continued)	Anilinopyrimidine	120	2.8	Supervised field trial	Individual compounds not specified	Kaethner, M., 2002
	Dicarboximide		5.6			
	Organophosphate		3.1			
	Pyrethroid		2.2			
	Triazole		2.3			
	Pyraclostrobin	120	1.8	Supervised field trial		Heck, W., 2002
Pyraclostrobin	120	2.5	Supervised field trial		Heck, W., 2002	
Pyraclostrobin	120	2.4	Supervised field trial		Heck, W., 2002	
Pyraclostrobin	120	2.4	Supervised field trial		Heck, W., 2002	
	Range (total mean)		1.8-3.4 (2.6)	Supervised field trial		
Potatoes	Chlorpropham	60	3.1	Supervised storage treatment		Quirijns, J.K., 2003
	Range (total mean)		3.1	Supervised storage treatment		
Head lettuce	Anilinopyrimidine	120	2.1	Supervised field trial	Individual compounds not specified	Kaethner, M., 2002a
	Triazole		2.0			
	Pyrethroid		1.9			
	Organophosphate		1.6			
	Dicarboximide		1.9			
	Carbamate	2.0				
	Anilinopyrimidine	120	2.0	Supervised field trial	Individual compounds not specified	Kaethner, M., 2002a
	Triazole		1.6			
	Pyrethroid		1.8			
Organophosphate		1.3				
Dicarboximide		1.8				
Carbamate		2.1				
Anilinopyrimidine	120	2.2	Supervised field trial	Individual compounds not specified	Kaethner, M., 2002a	
Triazole		1.8				
Pyrethroid		2.2				
Organophosphate		2.9				
Dicarboximide		2.5				
Carbamate		1.7				

Commodity	Pesticide	Number of units	Estimated variability factor	Type of survey	Remarks	Reference
Head lettuce (continued)	Anilinopyrimidine	120	1.3	Supervised field trial	compounds not specified	Kaethner, M., 2002a
	Triazole		1.3			
	Pyrethroid		1.3			
	Organophosphate		1.2			
	Dicarboximide		2.9			
	Carbamate		1.5			
	Range (total mean)		1.2-2.9 (1.9)	Supervised field trial		

A: Based on the ratio Maximum / Mean instead of 97.5 Percentile / Mean

B: Estimation of variability factor not possible, more than 50 % of the results were < LOQ

C: Market samples described as "unblended lots". However it is not clear, if this was really the case, so the data was not used to derive variability factors from